

# Ameliorative potential of vitamin E on the impact of dietary fumonisin B<sub>1</sub> on reproductive performance of female rabbits

Francis Ayodeji Gbore<sup>a,\*</sup>, Olufemi Adesanya Adu<sup>b</sup>

<sup>a</sup>Department of Animal Science, Adekunle Ajasin University, Akungba-Akoko, Nigeria

<sup>b</sup>Department of Animal Production and Health, The Federal University of Technology, Akure, Nigeria

## Abstract

Fumonisin B<sub>1</sub> (FB<sub>1</sub>), a contaminant of agricultural products, particularly maize worldwide is known to be consumed by farm animals and has been documented to cause various physiological responses in animals. A 15-week trial on the ameliorative potential of vitamin E on the negative impacts of FB<sub>1</sub> on reproductive performance of rabbits was conducted. Forty-nine female rabbits aged 16 to 18 weeks weighing 1.65 to 2.0 kg body weight were assigned to seven experimental feeding groups: the control group received a diet without FB<sub>1</sub>, three groups were fed diets containing different concentrations of FB<sub>1</sub> at 2.5, 5.0 or 7.5 mg kg<sup>-1</sup>, and three further groups had diets containing FB<sub>1</sub> and vitamin E i.e., 2.5 mg FB<sub>1</sub> kg<sup>-1</sup> + 100 mg vitamin E, 5.0 mg FB<sub>1</sub> kg<sup>-1</sup> + 100 mg vitamin E, and 7.5 mg FB<sub>1</sub> kg<sup>-1</sup> + 100 mg vitamin E. Data obtained on reproductive parameters - gestation length, litter size, kit weight as well as the kit crown-rump length, were analysed using ANOVA. Serum reproductive hormones - luteinizing hormone (LH), follicle stimulating hormone (FSH), prolactin, prostaglandin F<sub>2α</sub> (PGF<sub>2α</sub>), and estradiol (E<sub>2</sub>) levels in rabbits fed diets containing 7.5 mg FB<sub>1</sub> kg<sup>-1</sup> were significantly lower than those for all other treatments. Rabbits fed diets containing ≥ 5 mg FB<sub>1</sub> kg<sup>-1</sup> had significantly ( $p < 0.05$ ) longer gestation lengths and lower kit weights compared to the other treatment groups. The litter sizes of rabbits fed FB<sub>1</sub>-contaminated diets supplemented with vitamin E were significantly ( $p < 0.05$ ) higher compared with those on diets not supplemented, including the control. The 21-day *post-partum* weight gain of kits of does fed diets containing ≥ 5.0 mg FB<sub>1</sub> kg<sup>-1</sup> were significantly ( $p < 0.05$ ) lower than the weight gain observed in the other treatment groups. Does fed diets supplemented with vitamin E had significantly ( $p < 0.05$ ) higher milk yield compared with does on not supplemented diets, including the control. This study has shown that vitamin E supplementation of does counteracts the adverse impacts of FB<sub>1</sub> on reproductive hormones, gestation length, kit weight, and milk production in rabbits.

**Keywords:** rabbit, fumonisin B<sub>1</sub>, mycotoxin, reproduction, antioxidant

## 1 Introduction

Rabbit production has potential in many developing countries as a means of supplying cheap, high quality animal protein within the shortest time possible (Oyawaye *et al.*, 1990). Schiere (2004) stated that growth rate of around 15–20 g day<sup>-1</sup> are common in the tropics but it is possible to obtain 30–40 g day<sup>-1</sup> if rabbits are

well fed. Since good growth rate and reproductive performance form the basis of high profit margin in live-stock production, good nutrition is essential. However, some low-cost rabbit feed constituents, such as maize-milling waste, under humid tropical environmental conditions, may be infected with moulds and consequently may contain mycotoxins. Also, crops contaminated with high concentrations of mycotoxins are often diverted into animal feeds, thereby posing a serious threat to the growth, health and productivity of the animals (Griessler & Encarnaçao, 2009).

\* Corresponding author

Email: francis.gbore@aaua.edu.ng

Fumonisin B<sub>1</sub> (FB<sub>1</sub>), a mycotoxin produced by *Fusarium verticillioides* (= *F. moniliforme*) and other *Fusarium* species that grow on cereals, especially maize, has been documented to cause various physiological responses in humans and animals. The effects of dietary FB<sub>1</sub> on reproductive processes in animals have been well documented (Voss *et al.*, 2006; Gbore & Egbunike, 2008; Gbore, 2009a, b; Ewuola & Egbunike, 2010; Gbore *et al.*, 2012).

Considerable research has been directed at finding means to prevent or lessen the toxicity of FB<sub>1</sub>. The use of feed additives is an approach that is considered to be cheaper than degradation of the mycotoxin. Several studies have shown that a variety of adsorbent materials have a high affinity for binding mycotoxins by the formation of stable linkages (Huwig *et al.*, 2001; Galvano *et al.*, 2001; Diaz *et al.*, 2004; Var *et al.*, 2008). Experimental results obtained with some extensively studied adsorbents, such as hydrated sodium calcium aluminosilicate (HSCAS), are quite satisfactory with respect to aflatoxins, but they are not effective in preventing toxic effects of *Fusarium* mycotoxins, such as fumonisins, trichothecenes or zearalenone (Avantaggiato *et al.*, 2005). In the case of less- and non-absorbable mycotoxins such as these, new strategies have to be applied. Most of the mycotoxins provoke oxygen free radical formation (Balogh *et al.*, 2007; Pál *et al.*, 2009). As a result, the addition of natural or synthetic antioxidants in diets contaminated with these mycotoxins have been reported (Rogers, 2003; Cital *et al.*, 2005; Surai, 2006; Dvorska *et al.*, 2007) to be hypothetically effective in ameliorating the adverse impacts of these mycotoxins on animals due to their superoxide anion scavenging ability.

Dietary concentration of 7.5 mg fumonisin kg<sup>-1</sup> feed reportedly impaired reproductive performance in rabbits in 175 day studies (Ewuola & Egbunike, 2010). However, consumption of FB<sub>1</sub> at levels below this threshold may exert suboptimal reproductive performance in breeding female rabbits. The present study was therefore aimed at evaluating the ameliorative potential of vitamin E on the effects of dietary FB<sub>1</sub> below and above the no-observed-adverse-effect-level (NOAEL) of 5 mg kg<sup>-1</sup> feed on reproductive performance of female rabbits.

## 2 Materials and methods

### 2.1 Animals and experimental site

Forty-nine mixed breeds female rabbits aged 16 to 18 weeks were used in the current study. Rabbits were pro-

cured from a reputable commercial farm in Akure, Ondo State, Nigeria. This study was conducted in the Rabbit Unit of the Teaching and Research Farm of the Federal University of Technology, Akure, Nigeria. The farm is located in the humid rain forest zone of western Nigeria with mean rainfall, relative humidity and temperature of 1,500 mm, 75 % and 29 °C, respectively. The raining season is usually from March to November yearly. This study was carried out in accordance with “Guide for the care and use of Laboratory Animals” (NRC, 1996), and approved by the local Institutional Animal Ethics Committee.

### 2.2 Fumonisin B<sub>1</sub> production and experimental diets

Maize grits cultured with a toxigenic strain of *F. verticillioides* (MRC 286) and quantified in replicates for FB<sub>1</sub> and other common *Fusarium* mycotoxins as outlined by Gbore *et al.* (2016) were combined with non-contaminated maize grits to formulate seven diets consisted as follows: a control diet without FB<sub>1</sub> contamination, three diets containing different concentrations of FB<sub>1</sub> at 2.5, 5.0 or 7.5 mg FB<sub>1</sub> kg<sup>-1</sup> feed, and three further diets containing FB<sub>1</sub> and vitamin E at a fixed rate of 100 mg kg<sup>-1</sup> contaminated feed (i.e., 2.5 mg FB<sub>1</sub> kg<sup>-1</sup> + 100 mg vitamin E, 5.0 mg FB<sub>1</sub> kg<sup>-1</sup> + 100 mg vitamin E, and 7.5 mg FB<sub>1</sub> kg<sup>-1</sup> + 100 mg vitamin E). The level of FB<sub>1</sub> in the control diet was below the detection limit of 0.2 mg kg<sup>-1</sup> for the mycotoxin. The diets were marked A, B, C, D, E, F, and G, respectively, and provided nearly 15 % crude protein, 10 % crude fibre and 2600 kcal of digestible energy kg<sup>-1</sup> (Table 1).

**Table 1:** Ingredient and chemical composition (%) of the basal diet.

Ingredient	%
Maize †	40.7
Groundnut cake	11.8
Wheat offal	22.6
Palm kernel cake	19.6
Fish meal	2.0
Dicalcium phosphate	2.0
Salt (NaCl)	0.2
DL-Methionine	0.1
L-Lysine	1.06
Minerals/vitamins premix	0.1
<i>Analysed nutrients</i>	
Crude protein (%)	15.46
Crude fiber (%)	9.60
Digestible energy (kcal kg <sup>-1</sup> )	2597.96

† Varied proportion of *Fusarium*-contaminated and non-contaminated grains

### 2.3 Treatments and experimental layout

The rabbits were weighed at the end of a two week pre-experimental physiological adjustment period and randomly assigned to each of the diets ( $n = 7$  rabbits per treatment) and housed individually in hutches. The animals were fed with their respective experimental diets for 15 weeks.

### 2.4 Evaluation of reproductive performance of the does

After eight weeks of feeding the respective experimental diets, the rabbits were mated to intact bucks, at the ratio of one buck to two does. At parturition, gestation length, mean litter size, kit weight, and kit crown rump length were determined. The crown rump length of each of the kits was determined using thread calibrated in cm.

### 2.5 Determination of serum reproductive hormones

Serum level of each of luteinizing hormone (LH), follicle stimulating hormone (FSH), prolactin, prostaglandin  $F_{2\alpha}$  ( $PGF_{2\alpha}$ ), and estradiol ( $E_2$ ), was determined using appropriate test kits (Endocrines Technologies Inc., Newark, CA). All samples were run in triplicate in a single assay ( $n = 7$  per treatment).

### 2.6 Evaluation of post-partum performance

Rabbits were maintained on the respective diets through gestation till 21 days *post-partum* and the following parameters were determined: litter sizes, kit weights, and mortalities of the offspring at birth, 7, 14 and 21 days *post partum*. Milk yield was calculated using the formula proposed by Lebas *et al.* (1986) i. e., Milk production of does (g) = (Live weight new born at 21 days of age – Live weight of new born)\*1.18.

### 2.7 Statistical analysis

The design used for this experiment is Completely Randomized Design (CRD). Data obtained were subjected to statistical analysis using one-way ANOVA procedure of Statistical Analysis Systems (SAS, 2008). Duncan's multiple range tests of the same software was used to separate all means at 5% probability level. Results giving  $p$ -values of  $< 0.05$  were considered significantly different.

## 3 Results

### 3.1 Reproductive performance

Dietary  $FB_1$  generally influenced the concentrations of serum hormones examined (Table 2). The serum LH,

FSH, prolactin,  $PGF_{2\alpha}$ , and  $E_2$  levels in rabbits fed diets containing  $7.5 \text{ mg } FB_1 \text{ kg}^{-1}$  were significantly lower than those fed on the other treatment diets.

The reproductive performance of the does fed diets containing  $FB_1$  and vitamin E is shown in Table 3. The gestation length of the rabbits was significantly ( $p < 0.05$ ) influenced by the dietary treatments. Rabbits fed diets containing  $\geq 5 \text{ mg } FB_1 \text{ kg}^{-1}$  had significantly longer gestation period compared with does in other treatment groups. The litter sizes of rabbits fed  $FB_1$ -contaminated diets supplemented with vitamin E were significantly higher ( $p < 0.05$ ) compared with those on diets not supplemented with vitamin E, including the control. The total and relative weights of the kits of rabbits fed diets containing  $\geq 5.0 \text{ mg } FB_1 \text{ kg}^{-1}$  were significantly lower ( $p < 0.05$ ) than of those in other treatment groups. The crown-rump lengths of kits of rabbits fed diets containing  $2.5 \text{ mg } FB_1 \text{ kg}^{-1}$  were significantly ( $p < 0.05$ ) longer than those of the kits of rabbits in the control group and those on the diet containing  $5.0 \text{ mg } FB_1 \text{ kg}^{-1}$ .

### 3.2 Pre-weaning performance

Table 4 shows the pre-weaning performance of kits of does fed different diets. Weight gained by the kits was significantly ( $p < 0.05$ ) influenced by the dietary treatments. The mean weight of kits of does fed diets containing  $7.5 \text{ mg } FB_1 \text{ kg}^{-1}$  (Diet D) was significantly lower compared to those fed the control diet (Diet A), diets containing  $2.5 \text{ mg } FB_1 \text{ kg}^{-1}$  (Diet B) and  $\leq 5.0 \text{ mg } FB_1 \text{ kg}^{-1}$  supplemented with vitamin E (Diets E and F) at day 7 *post-partum*. At 21-day *post-partum*, weight gained by kits of does fed diets containing  $\geq 5.0 \text{ mg } FB_1 \text{ kg}^{-1}$  (Diets C and D) were significantly lower than the weights gained by kits of does in the other treatment groups, with the kits of does fed diets containing  $2.5 \text{ mg } FB_1 \text{ kg}^{-1}$  supplemented with vitamin E having the highest gain. The gain in weights of kits of does fed the control diet, diet containing  $2.5 \text{ mg } FB_1 \text{ kg}^{-1}$  and diets supplemented with vitamin E were significantly higher than those of kits of does fed diets containing  $5.0$  and  $7.5 \text{ mg } FB_1 \text{ kg}^{-1}$  without vitamin E supplementation.

The estimated milk yield of does fed diets containing  $FB_1$  or with vitamin E is shown in Fig. 1. Milk yield of the does was significantly ( $p < 0.05$ ) influenced by the dietary treatments. Does fed diets supplemented with vitamin E had significantly ( $p < 0.05$ ) higher milk yield compared with those on diets not supplemented, including the control. The milk yield of the does significantly decreased with increase in the concentrations of  $FB_1$  in the diets.

**Table 2:** Serum hormones of female rabbits fed diets containing FB<sub>1</sub> with or without vitamin E.

Parameters (ng ml <sup>-1</sup> )	Diet A	Diet B	Diet C	Diet D	Diet E	Diet F	Diet G	± SEM
	Control	2.5 mg FB <sub>1</sub>	5.0 mg FB <sub>1</sub>	7.5 mg FB <sub>1</sub>	2.5 mg FB <sub>1</sub> + Vitamin E	5.0 mg FB <sub>1</sub> + Vitamin E	7.5 mg FB <sub>1</sub> + Vitamin E	
Luteinizing hormone	5.23 <sup>a</sup>	5.20 <sup>a</sup>	5.20 <sup>a</sup>	4.53 <sup>b</sup>	5.00 <sup>a</sup>	5.27 <sup>a</sup>	5.00 <sup>a</sup>	0.15
Follicle stimulating hormone	17.37 <sup>a</sup>	17.80 <sup>a</sup>	17.00 <sup>a</sup>	14.20 <sup>b</sup>	17.00 <sup>a</sup>	17.23 <sup>a</sup>	17.93 <sup>a</sup>	0.69
Prolactin	2.03 <sup>a</sup>	2.00 <sup>a</sup>	1.90 <sup>a</sup>	1.53 <sup>b</sup>	2.10 <sup>a</sup>	2.03 <sup>a</sup>	2.07 <sup>a</sup>	0.15
Prostaglandin F <sub>2α</sub>	5.77 <sup>a</sup>	5.70 <sup>a</sup>	5.83 <sup>a</sup>	4.33 <sup>b</sup>	5.93 <sup>a</sup>	5.93 <sup>a</sup>	5.77 <sup>a</sup>	0.28
Estradiol	9.11 <sup>a</sup>	9.75 <sup>a</sup>	9.38 <sup>a</sup>	6.52 <sup>b</sup>	9.47 <sup>a</sup>	9.42 <sup>a</sup>	9.33 <sup>a</sup>	0.66

<sup>a,b</sup>: Means on the same row with different superscripts differ significantly ( $p < 0.05$ ).

**Table 3:** Reproductive performance of does fed diets containing FB<sub>1</sub> with or without vitamin E (Mean ± SE).

Parameters	Diet A	Diet B	Diet C	Diet D	Diet E	Diet F	Diet G
	Control	2.5 mg FB <sub>1</sub>	5.0 mg FB <sub>1</sub>	7.5 mg FB <sub>1</sub>	2.5 mg FB <sub>1</sub> + Vitamin E	5.0 mg FB <sub>1</sub> + Vitamin E	7.5 mg FB <sub>1</sub> + Vitamin E
Gestation length (d)	30.33±0.58 <sup>c</sup>	30.00±0.00 <sup>c</sup>	32.00±1.00 <sup>b</sup>	33.33±1.53 <sup>a</sup>	29.67±0.58 <sup>c</sup>	30.00±0.00 <sup>c</sup>	30.00±0.00 <sup>c</sup>
Litter size	4.67±1.15 <sup>ab</sup>	3.67±1.53 <sup>ab</sup>	3.67±0.58 <sup>ab</sup>	3.00±1.00 <sup>b</sup>	5.33±1.15 <sup>a</sup>	5.33±1.15 <sup>a</sup>	5.33±0.58 <sup>a</sup>
Litter weight (g)	224.80±57.43 <sup>ab</sup>	173.30±80.68 <sup>b</sup>	116.67±16.04 <sup>c</sup>	119.97±36.64 <sup>c</sup>	284.60±61.35 <sup>a</sup>	247.40±57.16 <sup>ab</sup>	235.57±11.05 <sup>ab</sup>
Kit weight (g)	45.83±4.01 <sup>ab</sup>	48.13±2.11 <sup>ab</sup>	46.23±5.40 <sup>ab</sup>	40.30±1.65 <sup>b</sup>	53.77±6.76 <sup>a</sup>	46.30±1.76 <sup>ab</sup>	53.83±4.27 <sup>a</sup>
Kit crown rump length (cm)	12.00±1.00 <sup>b</sup>	13.33±0.76 <sup>a</sup>	12.00±0.50 <sup>b</sup>	10.67±0.58 <sup>c</sup>	10.67±0.58 <sup>c</sup>	10.33±0.58 <sup>c</sup>	11.33±0.58 <sup>bc</sup>

<sup>a,b,c</sup>: Means on the same row with different superscripts differ significantly ( $p < 0.05$ ).

**Table 4:** Pre-weaning performance of kits of does fed diets containing FB<sub>1</sub> with or without vitamin E (Mean ± SE).


Parameters	Diet A	Diet B	Diet C	Diet D	Diet E	Diet F	Diet G
	Control	2.5 mg FB <sub>1</sub>	5.0 mg FB <sub>1</sub>	7.5 mg FB <sub>1</sub>	2.5 mg FB <sub>1</sub> + Vitamin E	5.0 mg FB <sub>1</sub> + Vitamin E	7.5 mg FB <sub>1</sub> + Vitamin E
Weight at birth (g)	45.83±4.01 <sup>ab</sup>	48.13±2.11 <sup>ab</sup>	46.23±5.40 <sup>ab</sup>	40.30±1.65 <sup>b</sup>	53.77±6.76 <sup>a</sup>	46.30±1.76 <sup>ab</sup>	53.83±4.27 <sup>a</sup>
Weight at day 7 (g)	83.53±6.76 <sup>ab</sup>	80.33±0.58 <sup>ab</sup>	77.80±7.12 <sup>bc</sup>	68.20±6.18 <sup>c</sup>	90.60±8.93 <sup>a</sup>	88.57±5.56 <sup>ab</sup>	78.83±2.76 <sup>bc</sup>
Weight at day 14 (g)	118.10±2.79 <sup>ab</sup>	117.87±2.58 <sup>ab</sup>	115.90±6.50 <sup>ab</sup>	99.40±11.46 <sup>c</sup>	127.27±5.99 <sup>a</sup>	123.67±4.56 <sup>a</sup>	120.40±7.24 <sup>bc</sup>
Weight at day 21 (g)	139.33±11.41 <sup>bc</sup>	135.00±5.00 <sup>c</sup>	116.27±7.11 <sup>d</sup>	114.07±8.93 <sup>d</sup>	155.30±3.12 <sup>a</sup>	142.43±3.82 <sup>ab</sup>	140.07±5.05 <sup>ab</sup>
Change in weight (g)	93.50±4.02 <sup>b</sup>	86.87±2.11 <sup>c</sup>	70.04±4.50 <sup>d</sup>	73.77±1.50 <sup>d</sup>	101.53±6.70 <sup>a</sup>	96.13±1.75 <sup>b</sup>	86.24±3.22 <sup>c</sup>
Mortality at Birth	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Mortality at day 7	1.00±1.00	1.33±0.58	0.33±0.58	1.33±0.58	0.33±0.58	1.00±0.00	0.33±0.58
Mortality at day 14	0.00±0.00	0.00±0.00	0.00±0.00	0.33±0.58	0.00±0.00	0.00±0.00	0.00±0.00
Mortality at day 21	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00

<sup>a,b,c,d</sup>: Means on the same row with different superscripts differ significantly ( $p < 0.05$ ).

## 4 Discussion

Although, previous studies (Voss *et al.*, 1996; LaBorde *et al.*, 1997; Collins *et al.*, 1998a,b) provided no evidence that FB<sub>1</sub> is a reproductive toxicant, more recent observations and experimental findings have however shown that FB<sub>1</sub> is a possible risk factor for birth defects (Merrill *et al.*, 2001; Marasas *et al.*, 2004; Voss *et al.*, 2006) and impaired reproductive capacity in animals (Gbore & Egbunike, 2008; Gbore, 2009a, b; Ewuola & Egbunike, 2010; Gbore *et al.*, 2012; Cortinovis *et al.*, 2014; Albonico *et al.*, 2016). In the present study, re-

sults show that > 5 mg dietary FB<sub>1</sub> kg<sup>-1</sup> feed had observable adverse effect on gestation length in rabbits. However, the inclusion of vitamin E at 100 mg kg<sup>-1</sup> diet countered this effect by reducing the gestation length. This could be attributed to the fact that vitamin E plays an important role in reproductive performance in animals due to its anti-oxidative properties which enabled its participation in metabolism of all cells as reported by Gutteridge & Halliwell (1994). Seemingly elongated gestation lengths was observed in rats fed *F. moniliforme* culture material containing 10 ppm FB<sub>1</sub> from two



**Fig. 1:** Mean milk yield of does.

weeks before mating by Voss *et al.* (1996). In a previous study, Gbore *et al.* (2012) observed significantly elongated gestation length from  $21.60 \pm 0.53$  days in rats fed diets containing  $\geq 10.0$  mg FB<sub>1</sub> kg<sup>-1</sup> to  $23.33 \pm 0.51$  days. These reports and the results from this study, further lead credibility to the fact that FB<sub>1</sub> potentially affects reproductive development in rabbits. The result of this study on gestation length shows that FB<sub>1</sub> at low concentration of  $2.5$  mg kg<sup>-1</sup> had no detrimental effect on the performance of rabbits but at higher concentrations of  $5.0$  and  $7.5$  mg kg<sup>-1</sup> the detrimental effects were observable.

Gbore *et al.* (2012) reported depressed serum gonadotropins levels in rats fed diets containing  $\geq 10$  mg FB<sub>1</sub> kg<sup>-1</sup>. The authors postulated that this might be due to an increase in gonadal steroid inhibition or suppression of the hypothalamus and/or pituitary gland with resultant decline in serum gonadotropins levels. Adverse effects of mycotoxins on sexual and reproductive developments have been reported. Dietary zearalenone levels as low as  $0.05$ – $0.06$  mg kg<sup>-1</sup> DM have been shown to increase the number of ovarian follicles and to decrease the serum concentration of the gonadotropic hormone (FSH) in female piglets (Döll *et al.*, 2003), thus potentially affecting their sexual development. Also, dietary FB<sub>1</sub> was reported to delay at-

tainment of sexual maturity in growing pigs (Gbore, 2009a) and rabbits (Ewuola & Egbunike, 2010). Fertility disturbances and other reproductive pathologies, particularly suppressive effect on testosterone secretion in mice, was reported (Yang *et al.*, 2007 a,b) following ingestion of cereals contaminated with *Fusarium* fungi. Also, reduced progesterone synthesis due to inhibition of the follicle stimulating hormone secretion (FSH) by *Fusarium* mycotoxins in cultured granulosa cells (GCs) from porcine ovaries was reported by Tiemann *et al.* (2003).

Improved litter size in rabbits fed diets supplemented with vitamin E over other treatment groups in this study may be attributed to the role vitamin E plays in reproduction. In the present study, it is evident that the dose-dependent significant decline in litter weight of does fed diets B, C, and D was countered by vitamin E supplementation. In a study, LaBorde *et al.* (1997) reported a reduced foetal weight from rabbits dosed daily by gavage on gestation days (GD) 3–19 with purified FB<sub>1</sub> at  $0.5$ – $1$  mg kg<sup>-1</sup> day<sup>-1</sup>. The authors ascribed this weight reduction to maternal toxicity, rather than any developmental toxicity produced by FB<sub>1</sub>. However, FB<sub>1</sub> dose-responsive significant decrease in foetal weight observed in this study correlates with reports of other studies. Pregnant rats dosed by gavage on GD 8–



12 with a semi-purified extract of culture material containing FB<sub>1</sub> with a purity of 80 % resulted in lower foetal weight at dose of 60 mg kg<sup>-1</sup> (Lebepe-Mazur *et al.*, 1995). Similarly, decreased body weight of live foetuses obtained from pregnant Syrian hamsters dosed with FB<sub>1</sub> in a dose-dependent manner was reported in a study by Penner *et al.* (1998). Voss *et al.* (1996) reported lower litter weights from rats fed *F. moniliforme* culture material providing 1–55 ppm FB<sub>1</sub> from two weeks before mating compared to the control group.

The significantly reduced litter weights at concentration of 7.5 mg FB<sub>1</sub> kg<sup>-1</sup> in this study corroborates the report of Voss *et al.* (1996) that FB<sub>1</sub> at high concentrations of 10 and 55 mg kg<sup>-1</sup> significantly reduced the weight of dams fed diets contaminated with *F. verticillioides*. It is also similar to the report of Lebepe-Mazur *et al.* (1995) that FB<sub>1</sub> at high dose significantly suppressed growth in rats. In the present study, vitamin E countered this effect by increasing the litter weight. The significant positive impact of vitamin E could be a result of the crucial role it plays in the growth of animals. It has been reported that FB<sub>1</sub> is not transferred through the placenta or into the milk in several animal species (Scott *et al.*, 1994; Becker *et al.*, 1995; Voss *et al.*, 1996; LaBorde *et al.*, 1997; Collins *et al.*, 1998b), nor are its metabolites found in animal products such as milk, meat and eggs (Jonker & van Egmond, 1999). Therefore, the resultant depressed growth rate of litters of does fed diets containing  $\geq 2.5$  mg FB<sub>1</sub> kg<sup>-1</sup> without vitamin E supplementation may be attributed to the reduced milk yields in does fed diets B, C, and D (Fig. 1).

Exposure to *Fusarium* mycotoxins has been linked to reproductive disorders in pigs (Cortinovic *et al.*, 2014; Gbore & Egbunike, 2008; Gbore, 2009a,b), mice (Yang *et al.*, 2007a,b), and rabbits (Ewuola & Egbunike, 2010). In a study to determine the potential reproductive effect of FB<sub>1</sub> on granulosa cell (GC) proliferation and steroid production in swine, Cortinovic *et al.* (2014) reported that this mycotoxin alone or with other *Fusarium* mycotoxins, including deoxynivalenol (DON) and zearalenone, influenced porcine GC proliferation and steroid production, thereby demonstrating their potential reproductive effects on swine. GC is reported to be crucial in the process of normal folliculogenesis and oocyte growth and development as they provide essential nutrients to the oocyte and establish a link between the oocyte and the surrounding ovarian tissue in which the follicle is embedded (Petro *et al.*, 2012). Also, GC is responsible for ovarian steroidogenesis that can be altered at any level leading to changes in the rate of hormone production and concentration (Ndossi *et al.*, 2012; Petro *et al.*, 2012). Progesterone is an essential regulator of

the reproductive events and plays key roles in ovulation, zygote implantation, and subsequent maintenance of pregnancy (Graham & Clarke, 1997). The influence of FB<sub>1</sub> on porcine GC function as noted by Cortinovic *et al.* (2014) could be one mechanism whereby this mycotoxin may impair reproductive activity in animals, particularly inhibiting effect on synthesis and secretion of gonadotropins. In rats, FB<sub>1</sub> concentrations at  $\geq 10$  mg kg<sup>-1</sup> diet were found to significantly reduce serum gonadotropin levels without inducing histopathological changes in the ovaries (Gbore *et al.*, 2012).

There exists a high correlation between the milk production by the doe and the growth of the kits because rabbit kits do not show significant feed intake before the age of 18–19 days (Fortun-Lamothe & Gidenne, 2000). Although Pascual *et al.* (1996) reported increase in milk yield with increase in litter sizes, the significantly decreased milk yield with increase in the concentrations of FB<sub>1</sub> in does fed diets supplemented with vitamin E, which had similar litter size, could be attributed to the mycotoxin in the diets. FB<sub>1</sub> bears a remarkable structural resemblance to sphinganine and sphingosine and the mycotoxin inhibits sphingosine metabolism in tissues, leading to an accumulation of sphingoid bases, which are intermediates in sphingolipid biosynthesis (Wang *et al.*, 1991). Sphingolipids are critical components for cell survival and homeostasis (Hirabayashi, 2012). High levels of sphingolipids have been reported in brains (Assi *et al.*, 2013) and its metabolism has a key neuropathological impact (Mencarelli & Martinez-Martinez, 2013). The disruption of sphingosine metabolism and alterations in the amounts of any of these intermediates by FB<sub>1</sub>, which could potentially alter neuronal function (Fonteh *et al.*, 2006) and result in a variety of biological effects (Penner *et al.*, 1998), may be responsible for the significant decline in serum gonadotropins and consequently reduced milk yield of does, with increased dietary FB<sub>1</sub> as observed in this study.

## 5 Conclusion

Economic loss in livestock production due to mycotoxins in animal feeds is a major problem in the tropics, including Nigeria. This study revealed that reproductive and pre-weaning performances of rabbits exposed to diets containing FB<sub>1</sub> could be impaired at concentrations above 5.0 mg kg<sup>-1</sup> of feed. Inclusion of vitamin E at 100 mg kg<sup>-1</sup> of feed in FB<sub>1</sub>-contaminated feeds is recommended as a nutritional strategy to alleviate the reproductive-depressing effect of the mycotoxin in female rabbits, especially at concentrations of  $\geq 5.0$  mg FB<sub>1</sub> kg<sup>-1</sup>.

## References

- Albonico, M., Schütz, L. F., Caloni, F., Cortinovis, C. & Spicer, L. J. (2016). Toxicological effects of fumonisin B<sub>1</sub> alone and in combination with other fusariotoxins on bovine granulosa cells. *Toxicon*, 118, 47–53.
- Assi, E., Cazzato, D., De Palma, C., Perrotta, C., Clementi, E. & Cervia, D. (2013). Sphingolipids and brain resident macrophages in neuroinflammation: An emerging aspect of nervous system pathology. *Clinical & Developmental Immunology*, 2013, Article ID 309302. doi:10.1155/2013/309302.
- Avantaggiato, G., Solfrizzo, M. & Visconti, A. (2005). Recent advances on the use of adsorbent materials for detoxification of *Fusarium* mycotoxins. *Food Additives and Contaminants*, 22, 379–388.
- Balogh, K., Hausenblasz, J., Weber, M., Erdélyi, M., Fodor, J. & Mézes, M. (2007). Effects of ochratoxin A on some production traits, lipid peroxide and glutathione redox status of weaned piglets. *Acta Veterinaria Hungarica*, 55, 463–470.
- Becker, B. A., Pace, L., Rottinghaus, G. E., Shelby, R., Misfeldt, M. & Ross, M. S. (1995). Effects of feeding fumonisin B<sub>1</sub> in lactating sows and their suckling pigs. *American Journal of Veterinary Research*, 56, 1253–1258.
- Citil, M., Gunes, V., Atakisi, O., Ozcan, A., Tuzcu, M. & Dogan, A. (2005). Protective effect of L-carnitine against oxidative damage caused by experimental chronic aflatoxicosis in quail (*Coturnix coturnix*). *Acta Veterinaria Hungarica*, 53, 319–324.
- Collins, T. F., Shackelford, M. E., Sprando, R. L., Black, T. N., LaBorde, J. B., Hansen, D. K., Eppley, R. M., Trucksess, M. W., Howard, P. C., Bryant, M. A., Ruggles, D. I., Olejnik, N. & Rorie, J. I. (1998a). Effects of fumonisin B<sub>1</sub> in pregnant rats. *Food and Chemical Toxicology*, 36, 397–408.
- Collins, T. F., Sprando, R. L., Black, T. N., Shackelford, M. E., LaBorde, J. B., Hansen, D. K., Eppley, R. M., Trucksess, M. W., Howard, P. C., Bryant, M. A., Ruggles, D. I., Olejnik, N. & Rorie, J. I. (1998b). Effects of fumonisins in pregnant rats, part 2. *Food and Chemical Toxicology*, 36, 673–685.
- Cortinovis, C., Caloni, F., Schreiber, N. B. & Spicer, L. J. (2014). Effects of fumonisin B<sub>1</sub> alone and combined with deoxynivalenol or zearalenone on porcine granulosa cell proliferation and steroid production. *Theriogenology*, 81, 1042–1049.
- Diaz, D. E., Hagler, W. M., Blackwelder, J. T., Eve, J. A., Hopkins, B. A., Anderson, K. L., Jones, F. T. & Whitlow, L. W. (2004). Aflatoxin binders II: Reduction of aflatoxin M1 in milk by sequestering agents of cows consuming aflatoxin in feed. *Mycopathologia*, 157, 233–241.
- Döll, S., Dänicke, S., Ueberschaer, K. H., Valenta, H., Schnurrbusch, U., Ganter, M., Klobasa, F. & Flachowsky, G. (2003). Effects of graded levels of *Fusarium* toxin contaminated maize in diets for female weaned piglets. *Archives of Animal Nutrition*, 57, 311–334.
- Dvorska, J. E., Pappas, A. C., Karadas, F., Speake, B. K. & Surai, P. F. (2007). Protective effect of modified glucomannans and organic selenium against antioxidant depletion in the chicken liver due to T-2 toxin-contaminated feed consumption. *Comparative Biochemistry and Physiology C Toxicology & Pharmacology*, 145, 582–587.
- Ewuola, E. O. & Egbunike, G. N. (2010). Effects of dietary fumonisin B<sub>1</sub> on the onset of puberty, semen quality, fertility rates and testicular morphology in male rabbits. *Reproduction*, 139, 439–445.
- Fonteh, A. N., Harrington, R. J., Huhmer, A. F., Biringer, R. G., Riggins, J. N. & Harrington, M. G. (2006). Identification of disease markers in human cerebrospinal fluid using lipidomic and proteomic methods. *Disease Markers*, 22, 39–64.
- Fortun-Lamothe, L. & Gidenne, T. (2000). The effects of size of suckled litter on intake behaviour, performance and health status of young and reproducing rabbits. *Annales de Zootechnie*, 49, 517–529.
- Galvano, F., Piva, A., Ritieni, A. & Galvano, G. (2001). Dietary strategies to counteract the effects of mycotoxins: a review. *Journal of Food Protection*, 64, 120–131.
- Gbore, F. A. (2009a). Growth performance and puberty attainment in growing pigs fed dietary fumonisin B<sub>1</sub>. *Journal of Animal Physiology and Animal Nutrition*, 93 (6), 761–767.
- Gbore, F. A. (2009b). Reproductive organ weights and semen quality of pubertal boars fed dietary fumonisin B<sub>1</sub>. *Animal*, 3 (8), 1133–1137.
- Gbore, F. A., Adu, O. A. & Ewuola, E. O. (2016). Protective role of supplemental vitamin E on brain acetylcholinesterase activities of rabbits fed diets contaminated with fumonisin B<sub>1</sub>. *European Journal of Biological Research*, 6 (2), 127–134.

- Gbore, F. A. & Egbunike, G. N. (2008). Testicular and epididymal sperm reserves and sperm production of pubertal boars fed dietary fumonisin B<sub>1</sub>. *Animal Reproduction Science*, 105, 392–397.
- Gbore, F. A., Owolawi, T. J., Erhunwunsee, M., Akele, O. & Gabriel-Ajobiwe, R. A. O. (2012). Evaluation of the reproductive toxicity of dietary fumonisin B<sub>1</sub> in rats. *Jordan Journal of Biological Science*, 5 (3), 183–190.
- Graham, J. D. & Clarke, C. L. (1997). Physiological action of progesterone in target tissues. *Endocrine Reviews*, 18, 502–519.
- Griessler, K. & Encarnaç o, P. (2009). Fumonisin – mycotoxins of increasing importance in fish! *Aquaculture Asia Magazine*, XIV (2), 24–26.
- Gutteridge, J. M. C. & Halliwell, B. C. (1994). Free radicals and antioxidants in ageing and diseases: Fact or fantasy. In: *Antioxidants in Nutrition, Health and Disease*. pp. 111–123, Oxford University Press, Oxford.
- Hirabayashi, Y. (2012). A world of sphingolipids and glycolipids in the brain -Novel functions of simple lipids modified with glucose. *Proceedings of the Japan Academy Series B Physical and Biological Sciences*, 88 (4), 129–143.
- Huwig, A., Freimund, S., Kappeli, O. & Dutler, H. (2001). Mycotoxin detoxification of animal feed by different adsorbents. *Toxicology Letters*, 122, 179–188.
- Jonker, M. A. & van Egmond, H. P. (1999). Mycotoxins in food of animal origin: A review. CRL document 389002095, CRL-European Union Community Reference Laboratory on Residues. RIVM-National Institute of Public Health and the Environment, Bilthoven, The Netherlands.
- LaBorde, J. B., Terry, K. K., Howard, P. C., Chen, J. J., Collins, F. X., Shackelford, M. E. & Hansen, D. K. (1997). Lack of embryotoxicity of fumonisin B<sub>1</sub> in New Zealand white rabbits. *Fundamental and Applied Toxicology*, 40, 120–128.
- Lebas, F., Coudert, P., Rouvier, R. & Rochambeau, H. D. (1986). The Rabbit husbandry, health and production. FAO Animal Production and Health Series, no. 21, FAO, Italy. Available at: <http://www.fao.org/docrep/x5082e/X5082E00.htm> (last accessed: 14.06.2017).
- Lebepe-Mazur, S., Bal, H., Hopmans, E., Murphy, P. & Hendrich, S. (1995). Fumonisin B<sub>1</sub> is fetotoxic in rats. *Veterinary and Human Toxicology*, 37, 126–130.
- Marasas, W. F., Riley, R. T., Hendricks, K. A., Stevens, V. L., Sadler, T. W., Gelineau-van Waes, J., Missmer, S. A., Cabrera, J., Torres, O., Gelderblom, W. C., Allegood, J., Mart nez, C., Maddox, J., Miller, J. D., Starr, L., Sullards, M. C., Roman, A. V., Voss, K. A., Wang, E. & Merrill Jr., A. H. (2004). Fumonisin disrupts sphingolipid metabolism, folate transport, and neural tube development in embryo culture and *in vivo*: a potential risk factor for human neural tube defects among populations consuming fumonisin-contaminated maize. *Journal of Nutrition*, 134, 711–716.
- Mencarelli, C. & Martinez-Martinez, P. (2013). Ceramide function in the brain: when a slight tilt is enough. *Cellular and Molecular Life Sciences*, 70 (2), 181–203.
- Merrill Jr., A. H., Sullards, M. C., Wang, E., Voss, K. A. & Riley, R. T. (2001). Sphingolipid metabolism: roles in signal transduction and disruption by fumonisin. *Environmental Health Perspectives*, 109 (Suppl 2), 283–289.
- Ndossi, D. G., Frizzell, C., Tremoen, N. H., F este, C. K., Verhaegen, S., Dahl, E., Eriksen, G. S., S rlie, M., Connolly, L. & Ropstad, E. (2012). An *in vitro* investigation of endocrine disrupting effects of trichothecenes deoxynivalenol (DON), T-2 and HT-2 toxins. *Toxicology Letters*, 214, 268–278.
- NRC (1996). *Guide for the Care and Use of Laboratory Animals*. National Research Council Institute of Laboratory Animal Resources (NRC), National Academy Press, Washington, DC.
- Oyawoye, E. O., Oyikin, M. E. & Shehu, Y. (1990). Studies in the nutrition of rabbits. Chemical evaluation of some tropical legumes as replacements for alfalfa in rabbit diets. *Journal of Applied Rabbit Research*, 13, 32–34.
- P l, L., Dublecz, K., Weber, M., Balogh, K., Erdelyi, M., Szigeti, G. & Mezes, M. (2009). Effect of Combined Treatment with Aflatoxin B-1 and T-2 Toxin and Metabolites on Some Production Traits and Lipid Peroxide Status Parameters of Broiler Chickens. *Acta Veterinaria Hungarica*, 57, 75–84.
- Pascual, J. J., Cervera, C., Blas, E. & Fern ndez-Carmona, J. (1996). Milk yield and composition in rabbit does using high fat diets. In: *Proceedings of the 6th World Rabbit Congress*, 9–12 July 1996, Toulouse, France, pp. 259–262. Vol. 1.



- Penner, J. D., Casteel, S. W., Pittman Jr, L., Rottinghaus, G. E. & Wyatt, R. D. (1998). Developmental toxicity of purified fumonisin B<sub>1</sub> in pregnant Syrian hamsters. *Journal of Applied Toxicology*, 18, 197–203.
- Petro, E. M., Leroy, J. L., Van Cruyten, S. J., Covaci, A., Jorssen, E. P. & Bols, P. E. (2012). Endocrine disruptors and female fertility: focus on (bovine) ovarian follicular physiology. *Theriogenology*, 78, 1887–1900.
- Rogers, S. A. (2003). Lipoic acid as a potential first agent for protection from mycotoxins and treatment of mycotoxicosis. *Archives of Environmental Health*, 58, 528–532.
- SAS (2008). *SAS/STAT User's Guide. Version 9.2 for Windows*. SAS Institute Inc., SAS Campus Drive. Cary, North Carolina, USA.
- Schiere, J. B. (2004). *Backyard Rabbit Farming in the Tropics*. Agromisa Foundation, Wageningen, the Netherlands. P. 42
- Scott, P. M., Delgado, T., Prelusky, D. B., Trenholm, H. L. & Miller, J. D. (1994). Determination of fumonisins in milk. *Journal of Environmental Science and Health Part B-Pesticides and Food Contaminants*, 29, 989–998.
- Surai, P. F. (2006). Selenium and mycotoxins. In: *Selenium in Nutrition and Health*, Ch. 6, pp. 317–362. Nottingham University Press, Nottingham.
- Tiemann, U., Tomek, W., Schneider, F. & Vanselow, J. (2003). Effects of the mycotoxins alpha- and beta-zearalenol on regulation of progesterone synthesis in cultured granulosa cells from porcine ovaries. *Reproductive Toxicology*, 17, 673–681.
- Var, I., Kabak, B. & Erginkaya, Z. (2008). Reduction in ochratoxin A levels in white wine, following treatment with activated carbon and sodium bentonite. *Food Control*, 19, 592–598.
- Voss, K. A., Bacon, C. W., Herbert, R. A., Chapin, R. E., Chamberlain, W. J., Plattner, R. D. & Meredith, F. I. (1996). Studies on the reproductive effects of *Fusarium moniliforme* culture material in rats and the biodistribution of [<sup>14</sup>C] fumonisin B<sub>1</sub> in pregnant rats. *Natural Toxins*, 4, 24–33.
- Voss, K. A., Gelineau-van Waes, J. B. & Riley, R. T. (2006). Fumonisin: current research trends in developmental toxicology. *Mycotoxin Research*, 22(1), 61–69.
- Wang, E., Norred, W. P., Bacon, C. W., Riley, R. T. & Merrill, A. H. (1991). Inhibition of sphingolipid biosynthesis by fumonisins: implications for diseases associated with *Fusarium moniliforme*. *Journal of Biological Chemistry*, 266, 14486–14492.
- Yang, Y. J., Wang, X. G., Liu, L. J. & Sheng, C. (2007a). Toxic effects of zearalenone and alpha zearalenol on the regulation of steroidogenesis and testosterone production in mouse Leydig cells. *Toxicology In Vitro*, 21(4), 558–565.
- Yang, Y. J., Wang, X. G., Liu, L. J. & Sheng, C. (2007b). Toxic effects of zearalenone and its derivatives zearalenol on male reproductive system in mice. *Reproductive Toxicology*, 24(3–4), 381–387.